Multiple Regression

Multicollinearity: Sleuth3 Chapter 12

Background

- Problems arise when too many explanatory variables are included in a model, particularly when some of these explanatory variables are correlated with each other
- In particular:
 - Precision in estimating important regression coefficients can be lost (meaning the variance is too high for those estimates to learn enough about the coefficients);
 - Prediction for a future response (value) may be negatively impacted
- The variance inflation factor (VIF) can help us measure the amount of multicollinearity in a set of candidate explanatory variables
 - Suppose you have an explanatory variable, X_j , in your regression model, and $R_{X_j}^2$ is the proportion of the variation in X_j that is explained by its relationship to other explanatory variables (this is like the coefficient of determination, R^2). Multicollinearity arises when X_j can be explained well by other explanatory variables. In other words, X_j is highly correlated with other explanatory variables in the model.
 - To determine the degree of multicollinearity between each X_j variable and the other explanatory variables in the model, we calculate:

$$VIF_j = \frac{1}{1 - R_{X_j}^2}.$$

VIF Rules of Thumb:

- VIF < 4: no multicollinearity between X_j and other explanatory variables
- $4 \leq VIF \leq 10$: moderate multicollinearity warrents further investigation
- VIF > 10: serious multicollinearity requires correction

Simulation with multicollinearity


Here I am going to simulate some data where some variables are correlated, and others are not to illustrate the multicollinearity issue and discuss cutoffs for VIF.

```
x1
                   x2
                             xЗ
## 2.369171 51.518287 24.420758 23.252615
confint(lm_sim)
                    2.5 %
                            97.5 %
##
## (Intercept) -1.9147608 2.6721559
              -0.5941507 2.2416647
## x1
## x2
              -0.2448766 2.8679931
## x3
              -0.7484275 1.3445816
## x4
              -1.6954309 0.3448929
```

Removing multicollinearity

Relationship to all subsets regression

```
candidate_models <- regsubsets(y ~ x1 + x2 + x3 + x4, data=sim_df)
plot(candidate_models)</pre>
```



summary(candidate_models)

```
## Subset selection object
## Call: regsubsets.formula(y ~ x1 + x2 + x3 + x4, data = sim_df)
## 4 Variables (and intercept)
     Forced in Forced out
##
## x1
         FALSE
                    FALSE
## x2
         FALSE
                    FALSE
## x3
         FALSE
                    FALSE
## x4
         FALSE
                    FALSE
## 1 subsets of each size up to 4
## Selection Algorithm: exhaustive
##
           x1 x2 x3 x4
## 1 (1) " " " " " * " "
## 2 (1) "*" "*" " "
## 3 (1) "*" "*" " "*"
## 4 ( 1 ) "*" "*" "*" "*"
```

summary(candidate_models)\$bic

```
## [1] -46.08163 -42.40569 -39.76786 -35.49870
```