# "Simple" Linear Regression

Sleuth 3: Chapter 7

### Example

We have a data set with information about 152 flights by Endeavour Airlines that departed from JFK airport in New York to either Nashville (BNA), Cincinnati (CVG), or Minneapolis-Saint Paul (MSP) in January 2012.

```
head(flights, 4)
## # A tibble: 4 x 3
##
     distance air_time dest
##
        <dbl>
                  <dbl> <chr>
## 1
         1029
                    189 MSP
## 2
          765
                    150 BNA
## 3
         1029
                    173 MSP
## 4
          589
                    118 CVG
```


#### So Far: ANOVA Model

- Observations in group i follow a Normal( $\mu_i$ ,  $\sigma^2$ ) distribution
- Observations are independent of each other

```
ggplot(data = flights, mapping = aes(x = air_time, color = dest)) +
geom_density() +
theme_bw()


0.06

dest
```



**Note:** The picture would look exactly the same if we treated distance as a categorical variable:

```
ggplot(data = flights, mapping = aes(x = air_time, color = factor(distance))) +
  geom_density() +
  theme_bw()
```




**Old idea:** Each group has a normal distribution with its own mean

• Categorical explanatory variable

New idea: Each group has a normal distribution with a mean that is a linear function of distance

• Quantitative (numeric) explanatory variable

## Warning: package 'ggridges' was built under R version 3.6.2



The simple linear regression is exactly like the ANOVA model, with the one new restriction that the means fall along a line.

#### Two ways to write the model:

#### Focusing on the mean (book)

Values of the response variable are independent and normally distributed with mean  $\mu(Y|X) = \beta_0 + \beta_1 X$ 

- Read as "The mean of Y for a given value of X"
- In our example, Y is air time and X is distance.

Written for a single observation, number i (my preference)

$$Y_i = \beta_0 + \beta_1 x_i + \varepsilon_i$$
  
 $\varepsilon_i \sim \text{Normal}(0, \sigma^2)$ 

• In our example,  $Y_i$  is the air time for flight number i and  $x_i$  is the distance for flight number i.

#### Parameter interpretations

- $\beta_0$  is intercept for the population: mean value of the response when X=0, in the population
- $\beta_1$  is slope for the population: change in mean response when X increases by 1 unit, in the population.
- $\beta_0$  and  $\beta_1$  are unknown population paramters. We estimate them with the intercept and slope of a line describing our sample.

## Conditions: spells "LINE-O"

Exactly the same as conditions for ANOVA, with addition that the mean of the response is a linear function of the explanatory varible:

- Linear relationship between explanatory and response variables
- **Independent** observations (knowing that one observation is above its mean wouldn't give you any information about whether or not another observation is above its mean)
- Normal distribution
- Equal standard deviation of response for all values of X
  - Denote this standard deviation by  $\sigma$
- no Outliers (not a formal part of the model, but important to check in practice)